


# 13: Differential Invariants & Proof Theory

## Logical Foundations of Cyber-Physical Systems




André Platzer

 Carnegie Mellon University  
Computer Science Department

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

- limits of computation
- proof theory for differential equations
- provability of differential equations
- nonprovability of differential equations
- proofs about proofs
- relativity theory of proofs
- inform differential invariant search
- intuition for differential equation proofs



core argumentative principles  
tame analytic complexity

improved analysis

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

## Differential Weakening

$$\frac{Q \vdash F}{P \vdash [x' = f(x) \& Q]F}$$

## Differential Invariant

$$\frac{Q \vdash [x' := f(x)](F)'}{F \vdash [x' = f(x) \& Q]F}$$


## Differential Cut

$$\frac{F \vdash [x' = f(x) \& Q]C \quad F \vdash [x' = f(x) \& Q \wedge C]F}{F \vdash [x' = f(x) \& Q]F}$$

DW  $[x' = f(x) \& Q]F \leftrightarrow [x' = f(x) \& Q](Q \rightarrow F)$

DI  $[x' = f(x) \& Q]F \leftarrow (Q \rightarrow F \wedge [x' = f(x) \& Q](F)')$

DC  $([x' = f(x) \& Q]F \leftrightarrow [x' = f(x) \& Q \wedge C]F) \leftarrow [x' = f(x) \& Q]C$



## Differential Weakening

$$\frac{Q \vdash F}{P \vdash [x' = f(x) \& Q]F}$$

## Differential Invariant

$$\frac{Q \vdash [x' := f(x)](F)'}{F \vdash [x' = f(x) \& Q]F}$$

## Differential Cut


$$\frac{F \vdash [x' = f(x) \& Q]C \quad F \vdash [x' = f(x) \& Q \wedge C]F}{F \vdash [x' = f(x) \& Q]F}$$

DW  $[x' = f(x) \& Q]F \leftrightarrow [x' = f(x) \& Q](Q \rightarrow F)$

DI  $[x' = f(x) \& Q]F \leftarrow (Q \rightarrow F \wedge [x' = f(x) \& Q](F)')$

DC  $([x' = f(x) \& Q]F \leftrightarrow [x' = f(x) \& Q \wedge C]F) \leftarrow [x' = f(x) \& Q]C$

DE  $[x' = f(x) \& Q]F \leftrightarrow [x' = f(x) \& Q][x' := f(x)]F$



- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

### Differential Invariant

$$\frac{Q \vdash [x' := f(x)](F)'}{F \vdash [x' = f(x) \& Q]F}$$

But generalizations are helpful to find the right  $F$  in the first place:

$$\text{cut,MR} \frac{A \vdash F \quad F \vdash [x' = f(x) \& Q]F \quad F \vdash B}{A \vdash [x' = f(x) \& Q]B}$$

### Compare Provability with Classes $\Omega$ of Differential Invariants

$\mathcal{DI}_\Omega$  : properties provable with differential invariants in  $\Omega \subseteq \{\geq, >, =, \wedge, \vee\}$

$\mathcal{A} \leq \mathcal{B}$  iff **all** properties provable with  $\mathcal{A}$  are also provable somehow with  $\mathcal{B}$

$\mathcal{A} \not\leq \mathcal{B}$  otherwise, i.e., **some** property can be proved with  $\mathcal{A}$  but not with  $\mathcal{B}$

$\mathcal{A} \equiv \mathcal{B}$  iff  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{B} \leq \mathcal{A}$  so **same** deductive power

$\mathcal{A} < \mathcal{B}$  iff  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{B} \not\leq \mathcal{A}$  so  $\mathcal{A}$  has strictly **less** deductive power

### Differential Invariant

$$\frac{Q \vdash [x' := f(x)](F)' \quad F \vdash [x' = f(x) \& Q]F}{F \vdash [x' = f(x) \& Q]F}$$

$\mathcal{DI}_{e=k} \equiv \mathcal{DI}_{e=0}$  by considering  $(e - k) = 0$

But generalizations are helpful to find the right  $F$  in the first place:

$$\text{cut,MR} \frac{A \vdash F \quad F \vdash [x' = f(x) \& Q]F \quad F \vdash B}{A \vdash [x' = f(x) \& Q]B}$$

### Compare Provability with Classes $\Omega$ of Differential Invariants

$\mathcal{DI}_\Omega$  : properties provable with differential invariants in  $\Omega \subseteq \{\geq, >, =, \wedge, \vee\}$

$\mathcal{A} \leq \mathcal{B}$  iff **all** properties provable with  $\mathcal{A}$  are also provable somehow with  $\mathcal{B}$

$\mathcal{A} \not\leq \mathcal{B}$  otherwise, i.e., **some** property can be proved with  $\mathcal{A}$  but not with  $\mathcal{B}$

$\mathcal{A} \equiv \mathcal{B}$  iff  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{B} \leq \mathcal{A}$  so **same** deductive power

$\mathcal{A} < \mathcal{B}$  iff  $\mathcal{A} \leq \mathcal{B}$  and  $\mathcal{B} \not\leq \mathcal{A}$  so  $\mathcal{A}$  has strictly **less** deductive power

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$F$  differential invariant of  $x' = f(x) \& Q$   
iff     $G$  differential invariant of  $x' = f(x) \& Q$

Proof.



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{l} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\text{MR,cut} \frac{}{F \vdash [x' = f(x) \& Q]F}$$



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\frac{\text{dl} \quad \frac{G \vdash [x' = f(x) \& Q]G}{}}{\text{MR,cut} \quad F \vdash [x' = f(x) \& Q]F}$$



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\begin{array}{c} [:=] \quad \overline{Q \vdash [x' := f(x)](G)'} \\ \text{dl} \quad \overline{G \vdash [x' = f(x) \& Q]G} \\ \text{MR,cut} \quad \overline{F \vdash [x' = f(x) \& Q]F} \end{array}$$



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\begin{array}{c} * \\ [=] \frac{}{Q \vdash [x' := f(x)](\textcolor{red}{F})'} \\ \text{dl} \quad \frac{}{G \vdash [x' = f(x) \& Q]G} \\ \text{MR,cut} \quad \frac{}{F \vdash [x' = f(x) \& Q]F} \end{array}$$



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\begin{array}{c} * \\ [=] \frac{}{Q \vdash [x' := f(x)](\textcolor{red}{F})'} \\ \text{dl} \quad \frac{G \vdash [x' = f(x) \& Q]G}{(F)' \leftrightarrow (G)' \text{ propositionally equivalent}} \\ \text{MR,cut} \quad F \vdash [x' = f(x) \& Q]F \end{array}$$



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is a propositional tautology then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\frac{\frac{\frac{*}{\frac{[::=] \frac{Q \vdash [x' := f(x)](\textcolor{red}{F})'}{G \vdash [x' = f(x) \& Q]G}}{\text{dl}}}{F \vdash [x' = f(x) \& Q]F}}{\text{MR,cut}}$$

$F \leftrightarrow G$  propositionally equivalent, so  
 $(F)' \leftrightarrow (G)'$  propositionally equivalent  
 since  $(F_1 \wedge F_2)' \equiv (F_1)' \wedge (F_2)'$  ...



Can use any propositional normal form

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$F$  differential invariant of  $x' = f(x) \& Q$   
iff     $G$  differential invariant of  $x' = f(x) \& Q$

Proof.



Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$F$  differential invariant of  $x' = f(x) \& Q$   
iff     $G$  differential invariant of  $x' = f(x) \& Q$

Proof.

$$\text{dI } \overline{-5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$



Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} & G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\frac{\text{:}=\overline{\vdash [x':=-x](0 \leq x' \wedge x' \leq 0)}}{\text{dl } \overline{-5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)}}$$



Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff} \quad G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\frac{\frac{\frac{\vdash 0 \leq -x \wedge -x \leq 0}{\vdash [x':=-x](0 \leq x' \wedge x' \leq 0)}}{\vdash [x' = -x](-5 \leq x \wedge x \leq 5)}}{\text{dl } -5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$



Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{c} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

not valid

$$\frac{}{\vdash 0 \leq -x \wedge -x \leq 0}$$

$$\frac{[:=]}{\vdash [x':=-x](0 \leq x' \wedge x' \leq 0)}$$

$$\frac{\text{dl}}{-5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$



Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{l} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

not valid

$$\vdash 0 \leq -x \wedge -x \leq 0$$

$$\vdash [x' := -x](0 \leq x' \wedge x' \leq 0)$$

$$[:=] \frac{}{-5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$

$$\text{dl } \frac{}{x^2 \leq 5^2 \vdash [x' = -x]x^2 \leq 5^2}$$

$$\text{arithmetic equivalence } -5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2$$

□

## Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{l} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

## Proof.

not valid

$$\vdash 0 \leq -x \wedge -x \leq 0$$

$$\frac{\vdash [x' := -x] (0 \leq x' \wedge x' \leq 0)}{\text{dl } \vdash [x' := -x] (-5 \leq x \wedge x \leq 5)}$$

$$\text{arithmetic equivalence } -5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2$$

$$\frac{\vdash [x' := -x] 2x x' \leq 0}{\text{dl } \vdash [x' := -x] x^2 \leq 5^2}$$

□

## Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{l} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

## Proof.

not valid

$$\vdash 0 \leq -x \wedge -x \leq 0$$

$$\vdash [x' := -x](0 \leq x' \wedge x' \leq 0)$$

$$\text{dl } \neg 5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5)$$

$$\text{arithmetic equivalence } -5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2$$

$$\mathbb{R} \quad \vdash \neg x 2x \leq 0$$

$$[:=] \quad \vdash [x' := \neg x] 2xx' \leq 0$$

$$\text{dl } x^2 \leq 5^2 \vdash [x' = -x] x^2 \leq 5^2$$

□

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is *real-arithmetic* equivalence then

$$\begin{array}{l} F \text{ differential invariant of } x' = f(x) \& Q \\ \text{iff } G \text{ differential invariant of } x' = f(x) \& Q \end{array}$$

Proof.

$$\frac{\text{not valid}}{\vdash 0 \leq -x \wedge -x \leq 0} \quad \frac{\mathbb{R} \quad *}{\vdash -x^2 \leq 0} \quad \frac{[:=]}{\vdash [x':=-x](0 \leq x' \wedge x' \leq 0)} \quad \frac{[:=]}{\vdash [x':=-x]2xx' \leq 0} \quad \frac{\text{dl} \quad \text{dl}}{\vdash -5 \leq x \wedge x \leq 5 \vdash [x' = -x](-5 \leq x \wedge x \leq 5) \quad \vdash x^2 \leq 5^2 \vdash [x' = -x]x^2 \leq 5^2}$$

$$\frac{\text{arithmetical equivalence } -5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2}{\square}$$

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is **real-arithmetical equivalence** then

$\exists$  differential invariant of  $x' = f(x) \wedge Q$   
 iff  $\exists$  differential invariant of  $x' = f(x) \wedge Q$

Proof.

not valid

$$\frac{\text{not valid}}{\vdash 0 \leq -x \wedge -x \leq 0} \quad \vdash [x' := -x](0 \leq x' \wedge x' \leq 0)$$

[:=]

$$\text{dl } \frac{-5 \leq x \wedge x \leq 5}{\vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$

$$\frac{*}{\vdash -x^2 \leq 0}$$







[:=]



$$\text{dl } \frac{\vdash [x' := -x]2xx' \leq 0}{\vdash x^2 \leq 5^2}$$

Despite arithmetic equivalence  $-5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2$













Differential structure matters! Higher degree helps here





Same  $p \geq 0$ .  
But different  $p' \geq 0$ .





Same  $p \geq 0$ .  
But different  $p' \geq 0$ .

Can still normalize  
atomic formulas to  
 $e = 0, e \geq 0, e > 0$

## Proposition (Equational deductive power [6, 2])

$$\mathcal{DI}_= \quad \mathcal{DI}_{=,\wedge,\vee}$$

Proof core.

Full: [6, 2].



Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$

Proof core.

Full: [6, 2].



Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2$
  
- $e_1 = e_2 \wedge k_1 = k_2$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$
- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee}$ 

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

□

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=\wedge,\vee}$

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

$$\text{So } [x' := f(x)](((e_1 - e_2)^2 + (k_1 - k_2)^2)' = 0)$$

$$\equiv [x' := f(x)](2(e_1 - e_2)((e_1)' - (e_2)') + 2(k_1 - k_2)((k_1)' - (k_2)')) = 0 \quad \square$$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=\wedge,\vee}$

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

So  $[x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

So  $[x' := f(x)](((e_1 - e_2)^2 + (k_1 - k_2)^2)' = 0)$

$$\equiv [x' := f(x)](2(e_1 - e_2)((e_1)' - (e_2)') + 2(k_1 - k_2)((k_1)' - (k_2)')) = 0 \quad \square$$

## Proposition (Equational deductive power [6, 2])

*atomic equations are enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=\wedge,\vee}$

Proof core.

Full: [6, 2].

- $e_1 = e_2 \vee k_1 = k_2 \leftrightarrow (e_1 - e_2)(k_1 - k_2) = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

So  $[x' := f(x)]((e_1 - e_2)(k_1 - k_2))' = 0$

$$\equiv [x' := f(x)](((e_1)' - (e_2)')(k_1 - k_2) + (e_1 - e_2)((k_1)' - (k_2)')) = 0$$

- $e_1 = e_2 \wedge k_1 = k_2 \leftrightarrow (e_1 - e_2)^2 + (k_1 - k_2)^2 = 0$

$$[x' := f(x)]((e_1)' = (e_2)' \wedge (k_1)' = (k_2)')$$

So  $[x' := f(x)](((e_1 - e_2)^2 + (k_1 - k_2)^2)' = 0)$

$$\equiv [x' := f(x)](2(e_1 - e_2)((e_1)' - (e_2)') + 2(k_1 - k_2)((k_1)' - (k_2)')) = 0 \quad \square$$

## Proposition (Equational [2])

$$\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} \quad \mathcal{DI} \quad \mathcal{DI}_\geq \quad \mathcal{DI}_\equiv$$

Proof core.



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$

$$\text{dl } \overline{x \geq 0 \vdash [x' = 5]x \geq 0}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$

$$\frac{\text{[:=]} \quad \vdash [x' := 5]x' \geq 0}{\text{dl } x \geq 0 \vdash [x' = 5]x \geq 0}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$

$$\frac{\mathbb{R} \quad \frac{}{\vdash 5 \geq 0}}{[:=] \quad \frac{}{\vdash [x':=5]x' \geq 0}} \text{ dl } \frac{}{x \geq 0 \vdash [x' = 5]x \geq 0}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$


$$\frac{\begin{array}{c} \mathbb{R} \quad \hline * \\ \hline \vdash 5 \geq 0 \end{array}}{[\!:=\!] \quad \hline \vdash [x':=5]x' \geq 0}$$
$$\text{dI } \frac{x \geq 0 \vdash [x' = 5]x \geq 0}{}$$



## Example (Sets Bijective or Not)

$$1 \longrightarrow 2 \longrightarrow 3 \longrightarrow 4 \longrightarrow 5 \longrightarrow 6$$
$$a \longrightarrow b \longrightarrow c \longrightarrow d \longrightarrow e \longrightarrow f$$


## Example (Vector Spaces Isomorphic or Not)



## Example (Sets Bijective or Not)

$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ | & & | & & | & & | & & | & & | \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & \longrightarrow & f \end{array}$$

## Example (Vector Spaces Isomorphic or Not)




## Example (Sets Bijective or Not)

$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ | & & | & & | & & | & & | & & | \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & \longrightarrow & f \end{array}$$

$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ & & & & & & & & & & \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & & \end{array}$$

## Example (Vector Spaces Isomorphic or Not)



## Example (Sets Bijective or Not)


$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ | & & | & & | & & | & & | & & | \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & \longrightarrow & f \end{array}$$

$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ & & & & & & & & & & \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & & \end{array}$$



criterion: cardinality  $|\{1, \dots, 6\}| = 6 \neq |\{a, b, c, d, e\}| = 5$

Need an indirect criterion especially if these sets are infinite

## Example (Vector Spaces Isomorphic or Not)




## Example (Sets Bijective or Not)





criterion: cardinality  $|\{1, \dots, 6\}| = 6 \neq |\{a, b, c, d, e\}| = 5$

Need an indirect criterion especially if these sets are infinite

## Example (Vector Spaces Isomorphic or Not)





### Example (Sets Bijective or Not)



criterion: cardinality  $|\{1, \dots, 6\}| = 6 \neq |\{a, b, c, d, e\}| = 5$

Need an indirect criterion especially if these sets are infinite

### Example (Vector Spaces Isomorphic or Not)



## Example (Sets Bijective or Not)



$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ | & & | & & | & & | & & | & & | \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & \longrightarrow & f \end{array}$$

$$\begin{array}{ccccccc} 1 & \longrightarrow & 2 & \longrightarrow & 3 & \longrightarrow & 4 & \longrightarrow & 5 & \longrightarrow & 6 \\ & & & & & & & & & & \\ a & \longrightarrow & b & \longrightarrow & c & \longrightarrow & d & \longrightarrow & e & & \end{array}$$

criterion: cardinality  $|\{1, \dots, 6\}| = 6 \neq |\{a, b, c, d, e\}| = 5$

Need an indirect criterion especially if these sets are infinite

## Example (Vector Spaces Isomorphic or Not)



criterion: dimension  $3 \neq 2$

## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$

$$\frac{\begin{array}{c} \mathbb{R} \quad \hline * \\ \hline \vdash 5 \geq 0 \end{array}}{[\!:=\!] \quad \hline \vdash [x':=5]x' \geq 0}$$
$$\text{dI } \frac{x \geq 0 \vdash [x' = 5]x \geq 0}{}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$       Unprovable with  $\mathcal{DI}_\equiv$

$$\frac{\text{R} \quad \frac{*}{\vdash 5 \geq 0}}{[:=] \quad \frac{\vdash [x':=5]x' \geq 0}{\text{dl} \quad \frac{x \geq 0 \vdash [x' = 5]x \geq 0}{\text{cut,MR} \quad \frac{}{x \geq 0 \vdash [x' = 5]x \geq 0}}}}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$ Unprovable with  $\mathcal{DI}_\equiv$ 

$$\frac{\mathbb{R} \quad \frac{*}{\vdash 5 \geq 0}}{[:=] \quad \frac{\vdash [x':=5]x' \geq 0}{\text{dl} \quad \frac{x \geq 0 \vdash [x' = 5]x \geq 0}{}}}$$

$$\frac{\text{dl} \quad \frac{p(x) = 0 \vdash [x' = 5]p(x) = 0}{\text{cut,MR} \quad \frac{}{x \geq 0 \vdash [x' = 5]x \geq 0}}}{}$$



Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_> \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_>$

## Unprovable with $\mathcal{DI}_\equiv$

$$\frac{\mathbb{R} \vdash 5 \geq 0}{\begin{array}{c} [:=] \vdash [x':=5]x' \geq 0 \\ \text{dI } x \geq 0 \vdash [x'=5]x \geq 0 \end{array}}$$

$$\frac{\text{dl} \quad \frac{}{p(x) = 0 \vdash [x' = 5]p(x) = 0}}{\text{cut,MR} \quad \frac{}{x \geq 0 \vdash [x' = 5]x \geq 0}}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$ 

$$\frac{\begin{array}{c} * \\ \mathbb{R} \quad \frac{}{\vdash 5 \geq 0} \\ \hline \vdash [x':=5]x' \geq 0 \end{array}}{[:=] \quad \frac{}{x \geq 0 \vdash [x' = 5]x' \geq 0}}$$

Unprovable with  $\mathcal{DI}_\equiv$ 

$$\frac{\begin{array}{c} ??? \\ \frac{}{\vdash [x':=5](p(x))' = 0} \\ \text{dl} \quad \frac{p(x) = 0 \vdash [x' = 5]p(x) = 0}{\text{cut,MR} \quad \frac{}{x \geq 0 \vdash [x' = 5]x' \geq 0}} \end{array}}{}$$



## Proposition (Equational incompleteness [2])

*Equations are not enough:*  $\mathcal{DI}_\equiv \equiv \mathcal{DI}_{=,\wedge,\vee} < \mathcal{DI}$  because  $\mathcal{DI}_\geq \not\leq \mathcal{DI}_\equiv$

Proof core.

Provable with  $\mathcal{DI}_\geq$ 

$$\frac{\begin{array}{c} * \\ \mathbb{R} \quad \frac{}{\vdash 5 \geq 0} \end{array}}{[\coloneqq] \quad \frac{}{\vdash [x':=5]x' \geq 0}} \quad \text{dl} \quad \frac{x \geq 0 \vdash [x' = 5]x \geq 0}{}$$

Unprovable with  $\mathcal{DI}_\equiv$ 

$$\frac{\begin{array}{c} ??? \\ \frac{}{\vdash [x':=5](p(x))' = 0} \\ \text{dl} \quad \frac{p(x) = 0 \vdash [x' = 5]p(x) = 0}{\text{cut,MR}} \end{array}}{x \geq 0 \vdash [x' = 5]x \geq 0}$$

Univariate polynomial  $p(x)$  is 0 if 0 on all  $x \geq 0$ 

Proposition (Strict barrier )

$\mathcal{DI}_>$     $\mathcal{DI}$     $\mathcal{DI}_=$     $\mathcal{DI}_>$

Proof core.



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.

Provable with  $\mathcal{DI}_=$

Unprovable with  $\mathcal{DI}_>$



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.

Provable with  $\mathcal{DI}_=$

Unprovable with  $\mathcal{DI}_>$

$$\text{dI } \overline{v^2 + w^2 = c^2 \vdash [v' = w, w' = -v] v^2 + w^2 = c^2}$$



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.

Provable with  $\mathcal{DI}_=$

Unprovable with  $\mathcal{DI}_>$

$$\frac{[::] \vdash [v' := w][w' := -v] 2vv' + 2ww' = 0}{\text{dl } v^2 + w^2 = c^2 \vdash [v' = w, w' = -v] v^2 + w^2 = c^2}$$



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.

Provable with  $\mathcal{DI}_=$

Unprovable with  $\mathcal{DI}_>$

$$\frac{\mathbb{R} \vdash 2vw + 2w(-v) = 0}{\begin{array}{l} [=] \quad \vdash [v':=w][w':=-v]2vv' + 2ww' = 0 \\ \text{dl } v^2 + w^2 = c^2 \vdash [v' = w, w' = -v]v^2 + w^2 = c^2 \end{array}}$$



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$*

Proof core.

Provable with  $\mathcal{DI}_=$   
\*

$$\frac{\mathbb{R} \vdash 2vw + 2w(-v) = 0}{\begin{array}{l} [=] \vdash [v' := w][w' := -v]2vv' + 2ww' = 0 \\ \text{dl } v^2 + w^2 = c^2 \vdash [v' = w, w' = -v]v^2 + w^2 = c^2 \end{array}}$$

Unprovable with  $\mathcal{DI}_>$



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.


Provable with  $\mathcal{DI}_=$

$$\frac{\begin{array}{c} \mathbb{R} \\ \vdash 2vw + 2w(-v) = 0 \\ \hline \text{[=:]} \quad \vdash [v' := w][w' := -v]2vv' + 2ww' = 0 \\ \text{dl } v^2 + w^2 = c^2 \vdash [v' = w, w' = -v]v^2 + w^2 = c^2 \end{array}}{*}$$


Unprovable with  $\mathcal{DI}_>$   
 $e > 0$  is open set.

$v^2 + w^2 = c^2$  is a closed set

closed  $v^2 + w^2 \leq 1$   
 with full boundary



open  $v^2 + w^2 < 1$   
 without boundary



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.


Provable with  $\mathcal{DI}_=$

$$\frac{\begin{array}{c} \mathbb{R} \\ \vdash 2vw + 2w(-v) = 0 \\ \hline \text{dl } v^2 + w^2 = c^2 \vdash [v' := w][w' := -v]2vv' + 2ww' = 0 \end{array}}{[v^2 + w^2 = c^2 \vdash [v' = w, w' = -v]v^2 + w^2 = c^2]}$$


Unprovable with  $\mathcal{DI}_>$   
 $e > 0$  is open set.  
 Only true and false  
 are both

$v^2 + w^2 = c^2$  is a closed set

closed  $v^2 + w^2 \leq 1$   
 with full boundary



open  $v^2 + w^2 < 1$   
 without boundary



## Proposition (Strict barrier incompleteness)

*Strict inequalities are not enough:*  $\mathcal{DI}_> < \mathcal{DI}$  because  $\mathcal{DI}_= \not\leq \mathcal{DI}_>$

Proof core.

Provable with  $\mathcal{DI}_=$

$$\frac{\begin{array}{c} \mathbb{R} \\ \hline \vdash 2vw + 2w(-v) = 0 \\ \hline \vdash [v' := w][w' := -v]2vv' + 2ww' = 0 \\ \hline \text{dl } v^2 + w^2 = c^2 \vdash [v' = w, w' = -v]v^2 + w^2 = c^2 \end{array}}{*}$$

Unprovable with  $\mathcal{DI}_>$

$e > 0$  is open set.


Only *true* and *false* are both

**but don't help proof**




$v^2 + w^2 = c^2$  is a closed set

closed  $v^2 + w^2 \leq 1$   
with full boundary



open  $v^2 + w^2 < 1$   
without boundary



Proposition (Equational )

$\mathcal{DI}_{=,\wedge,\vee}$      $\mathcal{DI}_{\geq}$

Proof core.



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.

Provable with  $\mathcal{DI}_{=}$

Provable with  $\mathcal{DI}_{\geq}$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.

Provable with  $\mathcal{DI}_{=}$ Provable with  $\mathcal{DI}_{\geq}$ 

$$\text{dI} \overline{e = 0 \vdash [x' = f(x) \& Q]e = 0}$$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.

Provable with  $\mathcal{DI}_{=}$

Provable with  $\mathcal{DI}_{\geq}$

$$\text{dI} \frac{\overline{Q \vdash [x' := f(x)](e)' = 0}}{e = 0 \vdash [x' = f(x) \& Q]e = 0}$$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.

Provable with  $\mathcal{DI}_{=}$

Provable with  $\mathcal{DI}_{\geq}$

$$\frac{\text{dI} \frac{*}{Q \vdash [x' := f(x)](e)' = 0}}{e = 0 \vdash [x' = f(x) \& Q]e = 0}$$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_{\geq}$

Proof core.

Provable with  $\mathcal{DI}_{=}$

Provable with  $\mathcal{DI}_{\geq}$

$$\frac{\frac{*}{Q \vdash [x' := f(x)](e)' = 0}}{\text{dI } e = 0 \vdash [x' = f(x) \& Q]e = 0}$$

$$\text{dI } -e^2 \geq 0 \vdash [x' = f(x) \& Q](-e^2 \geq 0)$$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_\geq$

Proof core.

Provable with  $\mathcal{DI}_=$ Provable with  $\mathcal{DI}_\geq$ 

$$\text{dI} \frac{\frac{*}{Q \vdash [x' := f(x)](e)' = 0}}{e = 0 \vdash [x' = f(x) \& Q]e = 0}$$

$$\text{dI} \frac{Q \vdash [x' := f(x)] - 2e(e)' \geq 0}{-e^2 \geq 0 \vdash [x' = f(x) \& Q](-e^2 \geq 0)}$$



## Proposition (Equational definability)

*Equations are definable by weak inequalities:*  $\mathcal{DI}_{=,\wedge,\vee} \leq \mathcal{DI}_\geq$

Proof core.

Provable with  $\mathcal{DI}_=$

$$\frac{\frac{*}{Q \vdash [x' := f(x)](\mathbf{e}') = 0}}{\text{dI } e = 0 \vdash [x' = f(x) \& Q]e = 0}$$

Provable with  $\mathcal{DI}_\geq$

$$\frac{\frac{*}{Q \vdash [x' := f(x)] - 2e(\mathbf{e}') \geq 0}}{\text{dI } -e^2 \geq 0 \vdash [x' = f(x) \& Q](-e^2 \geq 0)}$$



Local view of logic on differentials is crucial for this proof.

Degree increases

Theorem (Atomic

)

 $\mathcal{DI}_{\geq}$     $\mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>}$     $\mathcal{DI}_{>, \wedge, \vee}$ 

Proof idea.



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.

Provable with  $\mathcal{DI}_{\geq, \wedge, \vee}$

Unprovable with  $\mathcal{DI}_{\geq}$



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.

Provable with  $\mathcal{DI}_{\geq, \wedge, \vee}$

Unprovable with  $\mathcal{DI}_{\geq}$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \quad \vdash 5 \geq 0 \wedge y^2 \geq 0 \\
 \hline
 [:=] \quad \vdash [x' := 5][y' := y^2](x' \geq 0 \wedge y' \geq 0) \\
 \hline
 \text{dI } x \geq 0 \wedge y \geq 0 \vdash [x' = 5, y' = y^2](x \geq 0 \wedge y \geq 0)
 \end{array}$$



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.

Provable with  $\mathcal{DI}_{\geq, \wedge, \vee}$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \quad \frac{}{\vdash 5 \geq 0 \wedge y^2 \geq 0} \\
 \hline
 [:=] \quad \frac{}{\vdash [x':=5][y':=y^2](x' \geq 0 \wedge y' \geq 0)} \\
 \hline
 \text{dI} \quad \frac{}{x \geq 0 \wedge y \geq 0 \vdash [x' = 5, y' = y^2](x \geq 0 \wedge y \geq 0)}
 \end{array}$$

Unprovable with  $\mathcal{DI}_{\geq}$   
 $p(x, y) \geq 0 \leftrightarrow x \geq 0 \wedge y \geq 0$   
impossible since this implies  
 $p(x, 0) \geq 0 \leftrightarrow x \geq 0$   
so  $p(x, 0)$  is 0



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.

Provable with  $\mathcal{DI}_{\geq, \wedge, \vee}$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \quad \frac{}{\vdash 5 \geq 0 \wedge y^2 \geq 0} \\
 \hline
 [:=] \quad \frac{}{\vdash [x':=5][y':=y^2](x' \geq 0 \wedge y' \geq 0)} \\
 \hline
 \text{dI} \quad \frac{}{x \geq 0 \wedge y \geq 0 \vdash [x' = 5, y' = y^2](x \geq 0 \wedge y \geq 0)}
 \end{array}$$

Unprovable with  $\mathcal{DI}_{\geq}$   
 $p(x, y) \geq 0 \leftrightarrow x \geq 0 \wedge y \geq 0$   
impossible since this implies  
 $p(x, 0) \geq 0 \leftrightarrow x \geq 0$   
so  $p(x, 0)$  is 0

Substantial remaining parts of the proof shown elsewhere [2].



## Theorem (Atomic incompleteness)

*Atomic inequalities not enough:*  $\mathcal{DI}_{\geq} < \mathcal{DI}_{\geq, \wedge, \vee}$  and  $\mathcal{DI}_{>} < \mathcal{DI}_{>, \wedge, \vee}$

Proof idea.

Provable with  $\mathcal{DI}_{\geq, \wedge, \vee}$

$$\begin{array}{c}
 * \\
 \hline
 \mathbb{R} \quad \vdash 5 \geq 0 \wedge y^2 \geq 0 \\
 \hline
 [:=] \quad \vdash [x':=5][y':=y^2](x' \geq 0 \wedge y' \geq 0) \\
 \hline
 \text{dI } x \geq 0 \wedge y \geq 0 \vdash [x' = 5, y' = y^2](x \geq 0 \wedge y \geq 0)
 \end{array}$$

Unprovable with  $\mathcal{DI}_{\geq}$   
 $p(x, y) \geq 0 \leftrightarrow x \geq 0 \wedge y \geq 0$   
impossible since this implies  
 $p(x, 0) \geq 0 \leftrightarrow x \geq 0$   
so  $p(x, 0)$  is 0

Substantial remaining parts of the proof shown elsewhere [2]. □

dC still possible here but more involved argument separates.

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

Theorem (Gentzen's Cut Elimination)

(1935)

$$\frac{A \vdash B \vee C \quad A \wedge C \vdash B}{A \vdash B} \quad \text{cut can be eliminated}$$

Theorem (No Differential Cut Elimination)

(LMCS 2012)

*Deductive power with differential cuts exceeds deductive power without.*

$$\mathcal{DI} + \textcolor{red}{DC} > \mathcal{DI}$$

Theorem (Auxiliary Differential Variables)

(LMCS 2012)

*Deductive power with differential ghosts exceeds power without.*

$$\mathcal{DI} + DC + \textcolor{red}{DG} > \mathcal{DI} + DC$$

$$\text{dI } \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$\frac{[::=] \quad \vdash [x' := (x - 2)^4 + y^5][y' := y^2] 3x^2 x' \geq 0}{\text{dI } x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$\vdash 3x^2((x - 2)^4 + y^5) \geq 0$$

[:=]

$$\vdash [x' := (x - 2)^4 + y^5][y' := y^2]3x^2x' \geq 0$$

$$\text{dI } x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2]x^3 \geq -1$$

not valid

$$\vdash 3x^2((x - 2)^4 + y^5) \geq 0$$

[:=]

$$\vdash [x' := (x - 2)^4 + y^5][y' := y^2]3x^2x' \geq 0$$

$$\text{dI } x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2]x^3 \geq -1$$

not valid

$$\vdash 3x^2((x - 2)^4 + y^5) \geq 0$$

[:=]

$$\vdash [x' := (x - 2)^4 + y^5][y' := y^2]3x^2x' \geq 0$$

$$\text{dI } x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2]x^3 \geq -1$$

Have to know something about  $y^5$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$\text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] y^5 \geq 0}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$[:=] \frac{}{\vdash [x' := (x - 2)^4 + y^5][y' := y^2] 5y^4 y' \geq 0}$$
$$\text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] \textcolor{red}{y^5 \geq 0}}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

$$\begin{array}{c} \mathbb{R} \frac{}{\vdash 5y^4 y^2 \geq 0} \\ [=] \frac{}{\vdash [x' := (x - 2)^4 + y^5][y' := y^2] 5y^4 y' \geq 0} \\ \text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] y^5 \geq 0} \end{array}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] x^3 \geq -1}$$

\*

$$\mathbb{R} \frac{}{\vdash 5y^4y^2 \geq 0}$$

$$[:=] \frac{}{\vdash [x' := (x - 2)^4 + y^5][y' := y^2] 5y^4y' \geq 0}$$

$$\text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2] \textcolor{red}{y^5 \geq 0}}$$

$$\frac{\text{dI} \quad \overline{x^3 \geq -1 \vdash [x' = (x-2)^4 + y^5, y' = y^2 \& y^5 \geq 0] x^3 \geq -1 \triangleright}}{\text{dC} \quad \overline{x^3 \geq -1 \wedge \color{red}{y^5 \geq 0} \vdash [x' = (x-2)^4 + y^5, y' = y^2] x^3 \geq -1}}$$
$$\frac{*}{\mathbb{R} \quad \overline{\vdash 5y^4y^2 \geq 0}}$$
$$\frac{[:=] \quad \overline{\vdash [x':=(x-2)^4 + y^5][y':=y^2] 5y^4y' \geq 0}}{\text{dI} \quad \color{red}{y^5 \geq 0} \vdash [x' = (x-2)^4 + y^5, y' = y^2] \color{red}{y^5 \geq 0}}$$

$$[:=] \frac{}{y^5 \geq 0 \vdash [x' := (x - 2)^4 + y^5][y' := y^2]2x^2x' \geq 0}$$

$$\text{dI} \frac{}{x^3 \geq -1 \vdash [x' = (x - 2)^4 + y^5, y' = y^2 \& y^5 \geq 0]x^3 \geq -1 \triangleright}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2]x^3 \geq -1}$$

\*

$$\mathbb{R} \frac{}{\vdash 5y^4y^2 \geq 0}$$

$$[:=] \frac{}{\vdash [x' := (x - 2)^4 + y^5][y' := y^2]5y^4y' \geq 0}$$

$$\text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x - 2)^4 + y^5, y' = y^2]y^5 \geq 0}$$

$$\begin{array}{c}
 \hline
 \mathbb{R} \quad y^5 \geq 0 \vdash 2x^2((x-2)^4 + y^5) \geq 0 \\
 \hline
 [:=] \quad y^5 \geq 0 \vdash [x' := (x-2)^4 + y^5][y' := y^2]2x^2x' \geq 0 \\
 \hline
 \text{dI} \quad x^3 \geq -1 \vdash [x' = (x-2)^4 + y^5, y' = y^2 \& y^5 \geq 0]x^3 \geq -1 \triangleright \\
 \hline
 \text{dC} \quad x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x-2)^4 + y^5, y' = y^2]x^3 \geq -1
 \end{array}$$

\*

$$\begin{array}{c}
 \hline
 \mathbb{R} \quad \vdash 5y^4y^2 \geq 0 \\
 \hline
 [:=] \quad \vdash [x' := (x-2)^4 + y^5][y' := y^2]5y^4y' \geq 0 \\
 \hline
 \text{dI} \quad y^5 \geq 0 \vdash [x' = (x-2)^4 + y^5, y' = y^2]y^5 \geq 0
 \end{array}$$

\*

$$\mathbb{R} \frac{}{y^5 \geq 0 \vdash 2x^2((x-2)^4 + y^5) \geq 0}$$

$$[:=] \frac{}{y^5 \geq 0 \vdash [x' := (x-2)^4 + y^5][y' := y^2]2x^2x' \geq 0}$$

$$\text{dI} \frac{}{x^3 \geq -1 \vdash [x' = (x-2)^4 + y^5, y' = y^2 \& y^5 \geq 0]x^3 \geq -1 \triangleright}$$

$$\text{dC} \frac{}{x^3 \geq -1 \wedge y^5 \geq 0 \vdash [x' = (x-2)^4 + y^5, y' = y^2]x^3 \geq -1}$$

\*

$$\mathbb{R} \frac{}{\vdash 5y^4y^2 \geq 0}$$

$$[:=] \frac{}{\vdash [x' := (x-2)^4 + y^5][y' := y^2]5y^4y' \geq 0}$$

$$\text{dI} \frac{}{y^5 \geq 0 \vdash [x' = (x-2)^4 + y^5, y' = y^2]y^5 \geq 0}$$

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

Lemma (Differential invariants and propositional logic)

If  $F \leftrightarrow G$  is **real-arithmetical equivalence** then

$\exists$  differential invariant of  $x' = f(x) \wedge Q$   
 iff  $\exists$  differential invariant of  $x' = f(x) \wedge Q$

Proof.

not valid

$$\frac{\text{not valid}}{\vdash 0 \leq -x \wedge -x \leq 0} \quad \vdash [x' := -x](0 \leq x' \wedge x' \leq 0)$$

[:=]

$$\text{dl } \frac{-5 \leq x \wedge x \leq 5}{\vdash [x' = -x](-5 \leq x \wedge x \leq 5)}$$

$$\frac{*}{\vdash -x^2 \leq 0}$$

$\mathbb{R}$

[:=]

$$\frac{}{\vdash [x' := -x]2xx' \leq 0}$$

dl

$$\frac{x^2 \leq 5^2}{\vdash [x' = -x]x^2 \leq 5^2}$$

Despite arithmetic equivalence  $-5 \leq x \wedge x \leq 5 \leftrightarrow x^2 \leq 5^2$

□

Differential structure matters! Higher degree helps here

dC

---

$$A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_{\infty} \leq t$$

$$A \stackrel{\text{def}}{\equiv} v^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\|(x, y)\|_{\infty} \leq t \stackrel{\text{def}}{\equiv} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{\equiv} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

$$\frac{\text{dI} \quad \textcolor{red}{\square} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \textcolor{red}{v^2 + w^2 \leq 1}] \| (x, y) \|_{\infty} \leq t}{\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \| (x, y) \|_{\infty} \leq t}$$

$$A \stackrel{\text{def}}{=} v^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\| (x, y) \|_{\infty} \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\| (x, y) \|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

$$\frac{[:=] \overline{v^2 + w^2 \leq 1} \vdash [x' := v][y' := w][v' := \omega w][w' := -\omega v][t' := 1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')}{\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2 + w^2 \leq 1] \|(x, y)\|_\infty \leq t}$$

$$\frac{\text{dC}}{A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t}$$

$$A \stackrel{\text{def}}{\equiv} v^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{\equiv} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{\equiv} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

$$\mathbb{R} \frac{}{\nu^2 + w^2 \leq 1 \vdash -1 \leq \nu \leq 1 \wedge -1 \leq w \leq 1}$$

$$\frac{[:=] \frac{}{\nu^2 + w^2 \leq 1 \vdash [x' := \nu][y' := w][v' := \omega w][w' := -\omega v][t' := 1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')} {\text{dI} \quad \triangleleft \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \nu^2 + w^2 \leq 1] \|(x, y)\|_\infty \leq t} \text{dC} \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t$$

$$A \stackrel{\text{def}}{=} \nu^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\nu^2 + w^2 \leq 1 \vdash -1 \leq \nu \leq 1 \wedge -1 \leq w \leq 1}$$

$$\frac{[:=] \frac{}{\nu^2 + w^2 \leq 1 \vdash [x' := \nu][y' := w][v' := \omega w][w' := -\omega v][t' := 1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')} {\text{dI} \quad \triangleleft \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \nu^2 + w^2 \leq 1] \|(x, y)\|_{\infty} \leq t} \\ \text{dC} \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_{\infty} \leq t$$

$$A \stackrel{\text{def}}{\equiv} \nu^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\|(x, y)\|_{\infty} \leq t \stackrel{\text{def}}{\equiv} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{\equiv} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash -1 \leq v \leq 1 \wedge -1 \leq w \leq 1}$$

$$\frac{[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')} {\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \sqrt{v^2+w^2} \leq 1] \|(x, y)\|_{\infty} \leq t} \\ \text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_{\infty} \leq t$$

$$\text{dC} \frac{}{A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t}$$

$$A \stackrel{\text{def}}{=} \sqrt{v^2+w^2} \leq 1 \wedge x=y=t=0$$

$$\|(x, y)\|_{\infty} \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash -1 \leq v \leq 1 \wedge -1 \leq w \leq 1}$$

$$\frac{[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')} {\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \sqrt{v^2+w^2} \leq 1] \|(x, y)\|_\infty \leq t} \\ \text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t$$

$$\frac{\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \sqrt{v^2+w^2} \leq 1] \|(x, y)\|_2 \leq t}{\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t}$$

$$A \stackrel{\text{def}}{\equiv} \sqrt{v^2+w^2} \leq 1 \wedge x=y=t=0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{\equiv} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{\equiv} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash -1 \leq v \leq 1 \wedge -1 \leq w \leq 1}$$

$$\frac{[:=] \frac{}{v^2+w^2 \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')} \text{dI} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2+w^2 \leq 1] \|(x, y)\|_\infty \leq t}{\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t}$$

$$\frac{[:=] \frac{}{v^2+w^2 \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](2xv' + 2yw' \leq 2tt')} \text{dI} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2+w^2 \leq 1] \|(x, y)\|_2 \leq t}{\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t}$$

$$A \stackrel{\text{def}}{=} v^2+w^2 \leq 1 \wedge x=y=t=0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash -1 \leq v \leq 1 \wedge -1 \leq w \leq 1}$$

$$[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2 + w^2 \leq 1] \|(x, y)\|_\infty \leq t$$

$$\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t$$

$$\frac{}{\sqrt{v^2+w^2} \leq 1 \vdash 2xv + 2yw \leq 2t1}$$

$$[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](2xx' + 2yy' \leq 2tt')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2 + w^2 \leq 1] \|(x, y)\|_2 \leq t$$

$$\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t$$

$$A \stackrel{\text{def}}{=} \sqrt{v^2+w^2} \leq 1 \wedge x=y=t=0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\nu^2 + w^2 \leq 1 \vdash -1 \leq \nu \leq 1 \wedge -1 \leq w \leq 1}$$

$$[:=] \frac{}{\nu^2 + w^2 \leq 1 \vdash [x' := \nu][y' := w][v' := \omega w][w' := -\omega v][t' := 1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \nu^2 + w^2 \leq 1] \|(x, y)\|_\infty \leq t$$

$$\text{dC} \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t$$

not valid

$$\frac{}{\nu^2 + w^2 \leq 1 \vdash 2x\nu + 2yw \leq 2t}$$

$$[:=] \frac{}{\nu^2 + w^2 \leq 1 \vdash [x' := \nu][y' := w][v' := \omega w][w' := -\omega v][t' := 1](2xx' + 2yy' \leq 2tt')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& \nu^2 + w^2 \leq 1] \|(x, y)\|_2 \leq t$$

$$\text{dC} \quad A \vdash [x' = \nu, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t$$

$$A \stackrel{\text{def}}{=} \nu^2 + w^2 \leq 1 \wedge x = y = t = 0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$

$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

\*

$$\mathbb{R} \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash -1 \leq v \leq 1 \wedge -1 \leq w \leq 1}$$

$$[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](-t' \leq x' \leq t' \wedge -t' \leq y' \leq t')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2 + w^2 \leq 1] \|(x, y)\|_\infty \leq t$$

$$\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_\infty \leq t$$

Lower degree helps here

not valid

$$\frac{}{\sqrt{v^2+w^2} \leq 1 \vdash 2xv + 2yw \leq 2t}$$

$$[:=] \frac{}{\sqrt{v^2+w^2} \leq 1 \vdash [x':=v][y':=w][v':=\omega w][w':=-\omega v][t':=1](2xx' + 2yy' \leq 2tt')}$$

$$\text{dI} \quad \triangleleft \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1 \& v^2 + w^2 \leq 1] \|(x, y)\|_2 \leq t$$

$$\text{dC} \quad A \vdash [x' = v, y' = w, v' = \omega w, w' = -\omega v, t' = 1] \|(x, y)\|_2 \leq t$$

$$A \stackrel{\text{def}}{=} \sqrt{v^2+w^2} \leq 1 \wedge x=y=t=0$$

$$\|(x, y)\|_\infty \leq t \stackrel{\text{def}}{=} -t \leq x \leq t \wedge -t \leq y \leq t \quad \text{Supremum norm}$$


$$\|(x, y)\|_2 \leq t \stackrel{\text{def}}{=} x^2 + y^2 \leq t^2 \quad \text{Euclidean norm}$$

$$\forall x \forall y (\|(x, y)\|_{\infty} \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_{\infty})$$

$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_{\infty} \leq \|(x, y)\|_2 \right)$$


$$\forall x \forall y (\|(x, y)\|_\infty \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_\infty)$$

$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_\infty \leq \|(x, y)\|_2 \right)$$




$$\forall x \forall y (\|(x, y)\|_\infty \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_\infty)$$

$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_\infty \leq \|(x, y)\|_2 \right)$$



$$\forall x \forall y (\|(x, y)\|_\infty \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_\infty)$$

$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_\infty \leq \|(x, y)\|_2 \right)$$




$$\|\cdot\|_\infty \leq 1$$

$$\|\cdot\|_2 \leq 1$$

$$\|\cdot\|_\infty \leq \frac{1}{\sqrt{2}}$$

$$\forall x \forall y (\|(x, y)\|_\infty \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_\infty)$$

$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_\infty \leq \|(x, y)\|_2 \right)$$



$$\|\cdot\|_2 \leq \sqrt{2}$$


$$\|\cdot\|_\infty \leq 1$$

$$\|\cdot\|_2 \leq 1$$

$$\|\cdot\|_\infty \leq \frac{1}{\sqrt{2}}$$

$$\forall x \forall y (\|(x, y)\|_{\infty} \leq \|(x, y)\|_2 \leq \sqrt{n} \|(x, y)\|_{\infty})$$


$$\forall x \forall y \left( \frac{1}{\sqrt{n}} \|(x, y)\|_2 \leq \|(x, y)\|_{\infty} \leq \|(x, y)\|_2 \right)$$



Benefit from norm relations but be mindful of approximation error factors

- 1 Learning Objectives
- 2 Recap: Proofs for Differential Equations
- 3 Differential Equation Proof Theory
  - Propositional Equivalences
  - Differential Invariants & Arithmetic
  - Differential Structure
  - Differential Invariant Equations
  - Equational Incompleteness
  - Strict Differential Invariant Inequalities
  - Differential Invariant Equations to Differential Invariant Inequalities
  - Differential Invariant Atoms
- 4 Differential Cut Power & Differential Ghost Power
- 5 Curves Playing with Norms and Degrees
- 6 Summary

## Theorem (Differential Invariance Chart)



- Rich theory and structure behind differential invariants
- Scrutinize what property can be proved with what invariant
- Use provability sanity checks like open/closed/univariate
- Real differential semialgebraic geometry
- Exploit differential cuts to obtain more knowledge



André Platzer.

*Logical Foundations of Cyber-Physical Systems.*

Springer, Switzerland, 2018.

URL: <http://www.springer.com/978-3-319-63587-3>,

doi:10.1007/978-3-319-63588-0.



André Platzer.

The structure of differential invariants and differential cut elimination.

*Log. Meth. Comput. Sci.*, 8(4:16):1–38, 2012.

doi:10.2168/LMCS-8(4:16)2012.



André Platzer.

Foundations of cyber-physical systems.

Lecture Notes 15-424/624/824, Carnegie Mellon University, 2017.

URL: <http://lfcpss.org/course/fcps17.html>.



André Platzer.

A complete uniform substitution calculus for differential dynamic logic.

*J. Autom. Reas.*, 59(2):219–265, 2017.

doi:10.1007/s10817-016-9385-1.



André Platzer.

A differential operator approach to equational differential invariants.

In Lennart Beringer and Amy Felty, editors, *ITP*, volume 7406 of *LNCS*, pages 28–48, Berlin, 2012. Springer.

[doi:10.1007/978-3-642-32347-8\\_3](https://doi.org/10.1007/978-3-642-32347-8_3).



André Platzer.

Differential-algebraic dynamic logic for differential-algebraic programs.

*J. Log. Comput.*, 20(1):309–352, 2010.

[doi:10.1093/logcom/exn070](https://doi.org/10.1093/logcom/exn070).